


ABOUT THE AUTHOR



Parameswaran Nair, MD, PhD, FRCP, FRCPC

Dr. Nair graduated in 1989 from the University of Kerala Medical College in Trivandrum in India. He trained in Internal Medicine and in Respiratory Medicine at the University of Kerala, Royal Sussex County and Royal Sunderland Hospitals in the UK, and at McMaster University in Canada, where he did his doctoral thesis on leukotriene biology in asthma. He is currently the Frederick E. Hargreave Teva Innovation Chair in Airway Diseases & Professor of Medicine in the Division of Respiriology at McMaster University, an Adjunct Professor of Medicine at McGill University, and a Staff Respiriologist at the Firestone Institute for Respiratory Health at St. Joseph's Healthcare Hamilton. At the Firestone Institute, through inter-disciplinary clinics, he looks after patients with severe asthma and other complex airway and eosinophilic disorders, and provides access to them for biologics, molecular microbiology, novel pulmonary imaging, and bronchial thermoplasty, and opportunities to participate in research programs. His laboratory, funded by federal and provincial agencies, the Canada Research Chair program, AllerGen National Centre of Excellence, and industry partners, characterizes the types of bronchitis in airway diseases using biomarkers in sputum, identifies mechanisms of persistent bronchitis and airway hyperresponsiveness, and explores targeted therapies using small molecules and biologics. The patient-centred translational research program has been recognized by the Fellowships of the Canadian Academy of Health Sciences, the European Respiratory Society and the American College of Chest Physicians, the ATS Ann Woolcock Award, and the Asthma Society of Canada Bastable-Potts award, and has contributed to over 300 peer-reviewed publications (H-index 67, >21,000 citations) in major respiratory, allergy, and general medical journals. These observations, including 6 publications in the New England Journal of Medicine, paved the way for the development and approval of biologics that are currently available to treat patients with severe asthma.



Affiliations:

McMaster University
St. Joseph's Healthcare, Hamilton, Ontario

SEVERE ASTHMA PHENOTYPES: CHALLENGES AND OPTIONS FOR CONTROL

Introduction

The practice of medicine has evolved from “arm-chair medicine” through “evidence-based medicine” to “precision medicine”. Medical literature has seen a proliferation of the use of the phrases “precision medicine” and “personalized medicine,” with little distinction made between the two. While both strategies promote individualizing patient care, precision medicine is guided by information based on the genes, proteins, metabolites, and other biomarkers in the human body. In addition to these biological markers, personalized medicine would consider various social, economic, behavioural, and environmental factors that might be specific to a particular individual in planning a treatment strategy unique to that individual. The term P4 Medicine (Predictive, Preventive, Personalized and Participatory) has also been proposed to reflect the increased understanding and implications of the pathobiology of disease on management strategies.¹ The use of biologics and cell-based therapies, particularly in cancer therapeutics, has demonstrated the power of these strategies.² This brief review will focus on how this strategy is currently being applied in the management of severe asthma.

Definition of Phenotypes and Endotypes

A precise understanding of the pathobiology of disease and definitions is paramount to the practice of personalized medicine. The concept of “nominalism” as opposed to “essentialism” of asthma definitions is helpful to understand the mechanisms of disease and to identify the “treatable traits” that contribute to asthma symptoms and severity in individual patients.³ This involves identifying obvious clinical and physiological features (generally referred to as phenotypes),⁴ and specific biological characteristics that provide unique mechanistic insights (generally referred to as endotypes).⁵ These characteristics are generally identified based on unbiased hierarchical cluster analysis from data collected from large numbers of well-characterized patients in various cohort studies and cross-sectional studies.

Therefore, phenotypes include features such as: “early-onset or late-onset asthma”; “atopic or non-atopic asthma”; “obese or non-obese asthma”; “aspirin-sensitive or non-aspirin-sensitive asthma”; “smoker or non-smoker”; “nasal polyposis-associated

asthma or non-nasal polyposis-associated asthma”; and “asthma with fixed airflow obstruction or asthma with significant bronchodilator reversibility”. Triggers of bronchoconstriction, including occupational vs non-occupational; virus vs non-virus; exercise; cold air; and thunderstorm-related have also been employed to define phenotypes.⁶ While these clinical features are very helpful in predicting the clinical course of disease and, to some extent, responses to treatment, they do not provide precise information about the pathobiology of these features. A good example of this is so-called “exacerbation-prone asthma”. The reason for exacerbation might well be inappropriate therapy afforded to the patient rather than a patient characteristic.

Conversely endotyping endeavours to provide significant biological insights into specific disease or symptom manifestations. This requires the development and validation of simple clinically relevant and useful biomarkers that might reflect the underlying biology. The most widely employed marker is the eosinophil number in circulation or in the airway secretions (sputum). The term “eosinophilic” asthma is increasingly being used to classify patients with asthma whose symptoms and severity are believed to be mediated by the eosinophil, although this may not always be the case as a raised eosinophil number alone may not indicate that the key effector cell in the pathobiology of disease in that patient is the eosinophil. Thus, the presence or absence of eosinophils or their numbers is not an endotype. The mechanisms or cytokine pathways that lead to eosinophil recruitment are what truly constitute an endotype. The use of omics platforms has enabled the identification of genes (transcriptome); proteins (proteome); metabolites (metabolome); lipids (lipidome); and environmental factors interacting with biological factors (exposome), among other factors, to further endotype asthma.⁷

Identification of Phenotypes and Endotypes in Clinical Practice

A summary of the measurements or observations that are currently being used in our clinic at the Firestone Institute for Respiratory Health is shown in **Table 1**. Some of the examples of phenotyping guiding therapy include: “allergic asthma” responding well to allergen immunotherapy or to omalizumab; late-onset eosinophilic asthma responding well to anti-IL5

biologics; “aspirin-exacerbated asthma” responding to aspirin avoidance or aspirin desensitization; “obesity-related asthma” being associated with airway hyperresponsiveness and bronchomalacia; and “neutrophilic or non-eosinophilic asthma” characterized by susceptibility to recurrent airway infections.

Currently, there are only three biomarkers widely available for clinical use with which to gain insights into endotypes (although these markers are not precise indicators of the biology of the disease). These are total (or specific) IgE, eosinophil numbers in blood or in sputum, and the fraction of exhaled nitric oxide (FeNO). The relative merits and disadvantages of these three biomarkers have been extensively reviewed.⁸⁻¹⁰ While raised total IgE indicates allergy, it has very limited value in predicting response to a particular biologic. A combination of clinical features such as measuring airway hyperresponsiveness, FeNO, and blood (and preferably sputum) eosinophils can reasonably guide our choice of biologics for severe asthma. There are currently seven biologics approved for use in Canada.

Application of biomarkers and phenotyping/endotyping to manage severe asthma

A. Current clinical practice

Current clinical practice, which employs the above biomarkers, are endorsed by most National¹¹ and International¹² asthma guidelines including the Canadian Severe Asthma guidelines.¹³ The general principles are as follows:

- a) Confirm diagnosis of asthma with objective demonstration of variable airflow obstruction (peak flow, bronchodilator response or bronchoconstrictor response)
- b) Commence treatment with inhaled corticosteroids and bronchodilators (long-acting anti-cholinergics and long-acting beta-agonists), after checking inhaler technique and advising on allergen-avoidance measures, and regularly encouraging adherence to prescribed therapy. Allergen immunotherapy by adequately trained physicians is also effective in patients whose asthma is driven by one or two proven allergen sensitizations.¹⁴
- c) Most clinical trials have suggested that additional biomarkers are unlikely to make a significant difference to asthma outcomes compared to good clinical assessment and a spirometry.^{15,16}

- d) Increase the dose of inhaled corticosteroids and consider oral corticosteroid in patients with persistent raised blood eosinophil count (typically $>300/\mu\text{L}$). A persistently raised FeNO (typically >35 ppb), particularly when it is suppressed following a witnessed administration of corticosteroid, would indicate poor adherence.¹⁷
- e) With ongoing symptoms indicating poor control, additional endotyping/phenotyping is recommended. Early-onset asthma, young age and history of clinical allergies, along with raised blood eosinophil or total or specific IgE, would indicate omalizumab as the next step in the therapeutic algorithm. Conversely, late-onset asthma (whether associated with IgE or not); raised blood eosinophils, particularly associated with older age; nasal polyposis; and prednisone dependence (or frequent, 3-4 or more/year) might suggest an anti-IL5 biologic as the first choice.¹⁸ Currently available biomarkers do not help to differentiate between the three anti-IL5 biologics. A persistently raised FeNO (often >35 -50 ppb) despite normalizing or modest blood eosinophil count in patients with ongoing nasal polyposis or atopic dermatitis would suggest dupilumab as the next choice of biologic. Currently, there are no unique biomarkers that would help predict response to tezepelumab. It is claimed that biologics might be effective in all asthmatics including those with normal or low blood eosinophils or low FeNO. While this might be true for patients on high doses of inhaled corticosteroids, the efficacy of tezepelumab in the truly prednisone-dependent patients is still unproven.¹⁹

B. What can additionally be achieved in a research-supported environment?

The key to managing severe asthma and practicing precision medicine is to see patients at the time they are experiencing exacerbations,²⁰ (rather than adhering to the standard action plan of doubling inhaled steroids, or using prednisone or antibiotics); to assess if the symptoms (and reduction in airflow, FEV1) are due to luminal obstruction by eosinophils, neutrophils, other cell types, mucus, smooth muscle constriction, or airway wall thickness (or a combination of the above); and to target the dominant process that is contributing to the pathobiology (**Figure 1a**).²¹

A baseline blood eosinophil count, while a predictor of response to ALL biologics, does not help to discriminate between the biologics, and particularly is not helpful in monitoring the response to biologics or in assessing exacerbations while on biologics.²² Additional biomarkers that are (and can be) measured include: airway mucus (assessed by a CT mucus score);²³ sputum eosinophils and sputum neutrophils, sputum autoantibodies; sputum cytokines; and immunophenotyping. This includes assessment of primary and secondary immunodeficiencies (including whole exome sequencing), NK cell and macrophage functions, and the consequence of luminal obstruction by functional Xe¹²⁹ ventilation MRI. The application of these technologies to initiate and switch between biologics is summarized in **(Figure 1b)**.

This strategy²⁴ provides direction for deciding between the three anti-IL5 biologics considering the intensity of sputum eosinophilia, the presence of endogenous IgG autoantibodies in sputum, NK cell dysfunction, and anti-drug antibodies. For example, intense eosinophilia in sputum (often >20%) requiring >15 mg daily prednisone and in the presence of endogenous autoantibodies, would be associated with suboptimal response to the approved 100 mg subcutaneous dose of mepolizumab in almost 50% of prednisone-dependent patients.²⁵ Patients with NK cell dysfunction or anti-drug antibodies might demonstrate sub-optimal response to benralizumab.²⁶ Patients with severe airway hyper-responsiveness without demonstrable airway inflammation are likely to best respond to bronchial thermoplasty.²⁷ Persistent airway neutrophilia, unless otherwise proven, indicates an airway infection and should prompt investigations for susceptibility to such infections.²⁸

C. Future Diagnostic and Treatment Modalities

Airway measurements are critical in the management of severe asthma. Point of care assessments of airway eosinophilic²⁹ and neutrophilic activities³⁰ are likely to be available for clinical use. Comprehensive assessments, using large scale omics platforms, computational analytics and topological data analyses would precisely characterize the specific inflammatory pathways (Th2 high or Th2 low) and the

airway microbiome; and develop “handprints” for individual patients (as envisaged by the U-BIOPRED program),^{31,32} in order to truly individualize therapies for patients with severe asthma **(Figure 2)**.

Summary

In recent years tremendous advances in biotechnology have resulted in the development of effective medications, biologicals and technologies for bio-imaging and bio-intervention to treat patients with asthma. Currently, they are being more widely studied in adult asthmatics than in children with severe asthma. However, the same principles apply to the pediatric population.³³ What is required is to truly practice endotyping and precision medicine rather than giving perfunctory attention to this strategy. It is not possible to practice true precision medicine and to achieve optimal asthma control in every patient by solely monitoring a blood eosinophil count that is simply an overall marker of the burden of a Th2 disease. True personalized medicine requires: careful clinical examination; identifying phenotypes based on clinical characteristics; endotyping based on airway inflammatory responses and the pathways that modulate them; biomarkers and bio-imaging that reflect these processes; and selecting the appropriate therapy for the individual with consideration of their behavioural, social, economic, and environmental factors.

Correspondence:

Dr. Parameswaran Nair
Email: parames@mcmaster.ca

Financial Disclosures:

Grants to university: Sanofi, AZ, Teva, Roche, Genentech, Methapharm, Cyclomedica

Honoraria for lectures and Scientific Advisory Boards: AZ, Sanofi, GSK, Arrowhead

Phenotype	Investigations
CLINICAL	BLOOD
Atopy	CBC
Rhinosinusitis	Vasculitis workup
Obesity	Autoantibodies (blood/sputum)
Late-onset	B12, tryptase
ASA sensitivity	AAT
Infective bronchitis	Routine chemistry
PHYSIOLOGICAL	IMAGING/PULMONARY
Reversibility	CT/fMRI thorax
COPD	CT sinus
AHR	Echocardiogram
Tachyphylaxis	Full PFT
INFLAMMATORY	HAEMATOLOGY (when indicated)
Sputum eosinophilia	Bone marrow
Sputum neutrophilia	Cytogenetics
Blood eosinophilia	T-cell and receptors
Vasculitis	Cytokines
HES	
CURRENT TREATMENT	OTHER (as indicated)
Prednisone	GI (endoscopy)
High dose ICS	Stool
LABA	Skin prick test
LTRA	Bone density/Optomety
LAMA	EMG/NCV
Biologics	Sputum microbiome
Cytotoxics	FeNO
Antibiotics	
Nasal CS	CONSULTS
Sinus surgery	GI
Immunotherapy	Psychiatry
	ENT
ASTHMA EDUCATION	Allergy
Inhaler technique	Rheumatology
Compliance	Endocrinology
	Sleep medicine

Table 1 A checklist of biomarkers and endotyping to assess patients with severe asthma in an outpatient severe asthma clinic; courtesy of Parameswaran Nair, MD, PhD, FRCP, FRCPC




Figure 1a Contributors to severity and symptoms in asthma; reproduced with permission from Nair et al, 2021.




Figure 1b A strategy to initiate and switch biologics; reproduced with permission from Venegas Garrido C, et al, 2022




Figure 2. The concept for the future. Point of care tests would detect eosinophilic or neutrophilic activity in sputum that would then lead to the application of "omics" platforms to precisely identify the Th2 pathways or the microbial dysbiosis that leads to the cellular patterns, to guide therapy; courtesy of Parameswaran Nair, MD, PhD, FRCP, FRCPC

References

- Schüssler-Fiorenza Rose SM, Contrepolis K, Moneghetti KJ, Zhou W, Mishra T, Mataraso S, et al. A longitudinal big data approach for precision health. *Nat Med* 2019; 25:792–804.
- Hood L, Friend SH. Predictive, personalized, preventive, participatory (P4) cancer medicine. *Nature Rev Clin Oncol* 2011;8:184–87.
- Hargreave FE, Nair P. The definition and diagnosis of asthma. *Clin Exp Allergy* 2009; 39: 1652-8.
- Halder P, Pavord ID, Shaw DE, Berry MA, Thomas M, Brightling CE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med* 2008;178:218-224.
- Lötvall J, Akdis CA, Bacharier LB, Björner L, Casale TB, Custovic A, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011;127:355-60.
- Hekking PP, Bel EH. Developing and emerging clinical asthma phenotypes. *J Allergy Clin Immunol Pract* 2014;2:671-80.
- Wheelock CE, Goss VM, Balgoma D, Nicholas B, Brandsma J, Skipp PJ, et al. U-BIOPRED Study Group. Application of 'omics technologies' to biomarker discovery in inflammatory lung diseases. *Eur Respir J* 2013;42:802-25.
- Lim HF, Nair P. Airway Inflammation and Inflammatory Biomarkers. *Semin Respir Crit Care Med* 2018;39:56-63.
- Schleich FN, Seidel L, Sele J, Manise M, Quaedvlieg V, Michils A, et al. Exhaled nitric oxide thresholds associated with a sputum eosinophil count $\geq 3\%$ in a cohort of unselected patients with asthma. *Thorax* 2010;65:1039-44.
- Diamant Z, Vijverberg S, Alving K, Bakirtas A, Björner L, Custovic A, et al. Toward clinically applicable biomarkers for asthma: An EAACI position paper. *Allergy* 2019; 74: 1835-1851.
- National Institute for Health and Care Excellence. Asthma: diagnosis, monitoring and chronic asthma management <https://www.nice.org.uk/guidance/indevelopment/gid-ng10186/documents>, Accessed 28th February, 2023.
- Agache I, Akdis CA, Akdis M, Canonica GW, Casale T, Chivato T, et al. EAACI Biologicals Guidelines-Recommendations for severe asthma. *Allergy* 2021;76:14-44.
- FitzGerald JM, Lemiere C, Loughheed MD, et al. Recognition and management of severe asthma: A Canadian Thoracic Society position statement. *Can J Respir Crit Care Sleep Med*. 2017;1:199–221.
- Virchow JC, Backer V, Kuna P, Prieto L, Nolte H, Villesen HH, et al. Efficacy of a House Dust Mite Sublingual Allergen Immunotherapy Tablet in Adults With Allergic Asthma: A Randomized Clinical Trial. *JAMA* 2016;315:1715-25.
- Jayaram L, Pizzichini MM, Cook RJ, Boulet LP, Lemièrre C, Pizzichini E, et al. Determining asthma treatment by monitoring sputum cell counts: effect on exacerbations. *Eur Respir J* 2006;27:483-94.
- Calhoun WJ, Ameredes BT, King TS, Icitovic N, Bleecker ER, Castro M, et al. Comparison of physician-, biomarker-, and symptom-based strategies for adjustment of inhaled corticosteroid therapy in adults with asthma: the BASALT randomized controlled trial. Asthma Clinical Research Network of the National Heart, Lung, and Blood Institute. *JAMA* 2012;308:987-97.
- McNicholl DM, Stevenson M, McGarvey LP, Heaney LG. The utility of fractional exhaled nitric oxide suppression in the identification of nonadherence in difficult asthma. *Am J Respir Crit Care Med* 2012;186:1102-8.
- Rupani H, Murphy A, Bluer K, Renwick C, McQuitty P, Jackson DJ, et al. Biologics in severe asthma: Which one, When and Where? *Clin Exp Allergy* 2021;51:1225-8.
- Wechsler ME, Menzies-Gow A, Brightling CE, Kuna P, Korn S, Welte T, et al. Evaluation of the oral corticosteroid-sparing effect of tezepelumab in adults with oral corticosteroid-dependent asthma (SOURCE): a randomised, placebo-controlled, phase 3 study. *Lancet Respir Med* 2022; 10:650-60.
- Aziz-Ur-Rehman A, Dasgupta A, Kjarsgaard M, Hargreave FE, Nair P. Sputum cell counts to manage prednisone-dependent asthma: effects on FEV1 and eosinophilic exacerbations. *Allergy Asthma Clin Immunol* 2017; 13: 17. doi: 10.1186/s13223-017-0190-0.
- Nair P, O'Byrne PM. Medical algorithms: Approach to adult asthma exacerbations. *Allergy* 2021; 76:3556-9.
- Koenderman L, Hassani M, Mukherjee M, Nair P. Monitoring eosinophils to guide therapy with biologics in asthma: does the compartment matter? *Allergy* 2021;76: 1294-7.
- Svenningsen S, Haider E, Boylan C, Mukherjee M, Eddy RL, Capaldi DPI, et al. CT and Functional MRI to Evaluate Airway Mucus in Severe Asthma. *Chest* 2019;155:1178-9.
- Venegas Garrido C, Mukherjee M, Bhalla A, Nair P. Airway autoimmunity, asthma exacerbations, and response to biologics. *Clin Exp Allergy* 2022; 52:1365-8.
- Mukherjee M, Forero DF, Tran S, Boulay ME, Bertrand M, Bhalla A, et al. Suboptimal treatment response to anti-IL-5 monoclonal antibodies in severe eosinophilic asthmatics with airway autoimmune phenomena. *Eur Respir J* 2020;56:2000117. doi: 10.1183/13993003.00117-2020.
- Poznanski SM, Portillo A, Mukherjee M, Bhalla A, Son K, Ashkar AA, et al. Benralizumab's anti-eosinophil efficacy may be decreased by impaired NK cell activity. *Eur Respir J* 2022;59: 2102210. doi: 10.1183/13993003.02210-2021.
- Svenningsen S, Nair P, Eddy RL, McIntosh MJ, Kjarsgaard M, Lim HF, et al. Bronchial thermoplasty guided by hyperpolarised gas magnetic resonance imaging in adults with severe asthma: a 1-year pilot randomised trial. *ERJ Open Res* 2021;7: 00268-2021. doi: 10.1183/23120541.00268-2021.
- Nair P, Surette MG, Virchow JC. Neutrophilic asthma: misconception or misnomer? *Lancet Respir Med* 2021;9:441-3.
- Ali MM, Wolfe MG, Mukherjee M, Radford K, Patel Z, White D, et al. A sputum bioassay for airway eosinophilia using an eosinophil peroxidase aptamer. *Sci Rep* 2022;12:22476.
- Wolfe MG, Zhang Q, Hui C, Radford K, Nair P, Brennan JD. Development of a functional point-of-need diagnostic for myeloperoxidase detection to identify neutrophilic bronchitis. *Analyst* 2016; 141:6438-43.
- Ray A, Das J, Wenzel SE. Determining asthma endotypes and outcomes: Complementing existing clinical practice with modern machine learning. *Cell Rep Med* 2022;3:100857. Doi: 10.1016 / j.xcrm.2022.100857.
- Mazein A, Ivanova O, Balaur I, Ostaszewski M, Berzhitskaya V, Serebriyskaya T et al, AsthmaMap: An interactive knowledge repository for mechanisms of asthma. U-BIOPRED Study Group, eTRIKS Consortium. *J Allergy Clin Immunol* 2021;147:853-6.
- Bush A, Fitzpatrick AM, Saglani S, Anderson WC 3rd, Szeffler SJ. Difficult-to-Treat Asthma Management in School-Age Children. *J Allergy Clin Immunol Pract* 2022;10:359-75.